

Product Information

YF®532 Maleimide

Catalog Number: YM0049

Product Size: 1 mg

Application Scope: Fluorescent dye

Parameters

Ex/Em: 527/558 nm

Molecular Weight: ~810

Extinction Coefficient: 96,000

Substitute: Alexa Fluor 532 maleimide, CF 532 maleimide.

Storage

Store at -20°C and protect from light. When stored as directed, product is stable for at least 6 months.

Description

YF®532-Maleimide is a thiol reaction form of the YF® series of US Everbright dyes. Maleimide reacts with thiol groups to form thioether coupling products. The reaction can be carried out in an amine at pH = 7. In a neutral pH environment, the Maleimide group does not react with histidine or arginine.

Compared with other dyes (such as AlexaFluor®, DyLight®, Cy®Dyes or IRDyes®), YF® dyes are a new generation of fluorescent dyes with high brightness, light stability and water solubility advantages.

Protocol (Take labeled IgG antibody as an example)

1. Materials

- Anhydrous DMSO
- 10-100 mM phosphate (eg PBS), Tris or HEPES buffer, pH

7.0-7.5

- Cross-linked dextran G-25
- (Optional) Use Tris- (2-carboxyethyl) phosphine (TCEP) to reduce disulfide bonds in proteins to generate free sulfhydryl groups.
- NaN₃
- BSA

2. Labeled methods and steps

2.1 Prepare labeled antibodies

a) Dissolve the antibody in buffer at room temperature to a final concentration of 50-100 μM (7.5-15 mg / mL for IgG).

b) Optional step: If you want to release more thiol groups from the disulfide bonds in the protein, you can add about 10-fold molar excess of TCEP at this stage. Incubate the reaction solution for about 30 min. The reduction reaction and subsequent labeling reaction are preferably performed in the presence of an inert gas (N₂ or Ar) to prevent the re-formation of disulfide bonds.

2.2 Prepare Dye Storage Solution

Remove the YF®532-Maleimide dye and return to room temperature. Prepare a 10 mM dye stock solution. For 1 μmol dye: Add 100 μL of anhydrous DMSO to the vial. Vortex briefly to completely dissolve the dye, and then centrifuge for a short time to concentrate the dye on the bottom of the tube.

Note: 1) If a small amount of protein is used for the labeling reaction, the dye stock solution may need to be diluted to a lower concentration to ensure the accuracy of the volume



measurement.

2) The unused storage solution can be stored in a dry place at -20 °C, protected from light, for subsequent use. If anhydrous DMSO is used to prepare the solution, the dye is stable for at least one month.

3) Dye stock solution can also be prepared with dH₂O or aqueous buffer. However, because the dye will hydrolyze over time, the water-soluble storage solution needs to be ready-to-use and cannot be stored and used again.

2.3 Labeling reaction steps

a) While stirring or vortexing the protein solution, add a certain amount of dye stock solution to produce a mixed solution of YF[®] dye and protein molar ratio of 10-20. For example, for 50 μM IgG, the dye should be added to a final dye concentration of 0.5-1 mM.

b) Stir the reaction at room temperature in the dark for 2 h or overnight at 4 °C.

Note: While performing the labeling reaction, proceed to step 2.4a to prepare the glucan column.

2.4 Isolation of labeled proteins from the reaction solution

a) Equilibrate a dextran column (10 mm × 300 mm) with PBS buffer (pH ~ 7.4).

b) Load the reaction solution from step 2.3b onto the column and elute the column with PBS buffer. The first band eluting from the column corresponds to the antibody conjugate.

Note: For small-scale labeling reactions, to avoid excessive dilution of the product, an ultrafiltration device can be used to remove free dye in the conjugate. 10K ultrafiltration tubes can be used for IgG proteins; proteins with different molecular weights require different ultrafiltration tubes.

3. Determine the DOL

3.1 Determination of protein concentration

The antibody concentration can be calculated by the following

formula: $C \text{ (mg / mL)} = ((A_{280} - (A_{\text{max}} \times C_f)) / 1.4) \times \text{dilution factor}$.

■ C refers to the antibody concentration collected in the experiment;

■ Dilution factor refers to the dilution factor during photometric measurement;

■ A₂₈₀ and A_{max} refer to the absorbance at 280 nm and the absorbance at the absorption wavelength (~532 nm), respectively;

■ C_f is the correction factor, YF[®]532-Maleimide has a C_f value of 0.06;

■ 1.4 refers to the extinction coefficient of IgG (mL / mg);

Note: The protein solution eluted through the column is directly used for absorbance detection. The concentration may be too large, so it needs to be diluted to about 0.1 mg / mL. The dilution factor (dilution factor) needs to be estimated from the initial amount of antibody (eg 5 mg) and the total volume of protein solution eluted.

3.2 DOL estimation

DOL is calculated by the following formula: $\text{DOL} = (A_{\text{max}} \times \text{Mwt} \times \text{dilution factor}) / (\epsilon \times C)$

■ A_{max}, dilution factor, C value has been specified in 3.1;

■ Mwt refers to the molecular weight of IgG (~ 150,000);

■ ε is the molar absorption coefficient of YF[®]532-Maleimide;

Notes

1. When the protein labeled by this product needs to be stored for a long time, it is recommended to add 5-10 mg / mL BSA and 0.01-0.03% NaN₃ to prevent protein denaturation and microbial growth.

2. Store at 4 °C in the dark. If glycerin is added to a final concentration of 50%, it can be stored at -20 °C. Stable for more than one year.

